


Mal de Debarquement Survey, 2007

- Total number of responses= 101
- 96 responses with positive answers to Questions 1-3
- Survey conducted at an online site with one response per person being controlled.
- All slides after the second one are based only on those who responded yes to questions 1-3.



Questions One-Three



Do you feel like you are rocking, swaying or bobbing?


Does the feeling decrease when you are in motion such as in a car or boat?

Have your symptoms been persistent for at least 30 days?



Question 4


If you have undergone medical testing, were you tested for superior canal dehiscence?



22- Yes 53- NO

Question 5


If you have undergone medical testing, were you tested for perilymphatic fistula?




24- Yes 55- No



Question Six- Age at Onset



Total number of responses = 96




10 Male

86 Female



Question 8

What triggered persistent rocking (30 days or more)?



- A) Boat travel 3 days or less
- B) Boat travel 4 days or more
- C) Airline travel 4 hours or less
- D) Airline travel more than 4 hours
- E) Train travel 3 days or less
- F) Train travel more than 3 days
- G) Car travel 3 hours or less
- H) Car travel more than 3 hours
- I) Other* See next slide

Responders were able to choose more than one answer when applicable as many had combined modes of travel in the one episode that started the persistent symptoms.



Question 8 - Other Included

- 5- Elevator Use
- 2- Amusement Park Rides
- 4- Flu-like symptoms
- 3- Spontaneous
- Misc.- waterbed, unstable trailer type building, exercise equipment, combination of travel types.




Question 9- Repeat Episodes

- Thirty people responded that they have had repeat episodes from plane, boat, car, and train travel. Others reported repeat episodes from a motorcycle trip, tipping head when cleaning, RV travel, illness, surgery, stress, and tiredness.



Question 10

What is the duration of your rocking?



96 Responders

A) Continuous

B) Continuous except when I am in motion (walking, riding in a vehicle, etc.)

C) 12 hours or less per day

D) 6 hours or less per day

E) 1 hour or less per day


F) not daily



[MdDS Balance Disorder Foundation](http://www.mddsfoundation.org) May, 2007

www.mddsfoundation.org

Who Made the Diagnosis?




Some people received diagnoses from more than one practitioner.

Other responses included a Psychiatrist, Head of a Balance Center and a Neurological Chiropractor.



[MdDS Balance Disorder Foundation](http://www.mddsfoundation.org) May, 2007

www.mddsfoundation.org



Other included: Time, Inderal, combination of exercise and medication, lots of sleep, Chiropractic, massage therapy, and laying back in recliner.

